

Dynamic Optimization

Jesús Bueren

EUI

Dynamic Optimization

- In this chapter we are going to characterize solutions to dynamic optimization problems
- In order to solve them, we are going to introduce discrete dynamic programming.
- Along our way, we are going to revise some mathematical concepts covered by Villanacci.
- References: *The PhD Macro Book* (Ch 4), Acemoglu (Ch 6), and SLP (Ch 4).

Motivating the Recursive Formulation

A Cake Eating Problem

- We will go over a very simple dynamic optimization problem.
- Suppose that you are presented with a cake of size W_1 .
- At each point in time $t = 1, 2, \dots, T$, you can eat some of the cake but must save the rest.
- Let c_t be your consumption at time t and $u(c_t)$ represent the flow of utility.
- u twice differentiable, strictly increasing, strictly concave,
 $\lim_{c \rightarrow 0} u'(c) = \infty$.
- Discount factor: $0 < \beta < 1$

The Sequential Formulation

A Cake Eating Problem

- The agent is solving:

$$\begin{aligned} \max_{\{c_t, W_{t+1}\}_{t=0}^T} & \sum_{t=0}^T \beta^t u(c_t) \\ \text{s.t.} & c_t + W_{t+1} = W_t \quad \forall t \\ & W_{T+1} \geq 0 \end{aligned}$$

- The Lagrangian associated to this problem is given by:

$$\mathcal{L} = \sum_{t=0}^T \beta^t u(c_t) + \sum_{t=0}^T \lambda_t (W_t - c_t - W_{t+1}) + \phi W_{T+1}$$

The Sequential Formulation

A Cake Eating Problem

- FOCs:

$$\beta^t u_c(c_t) = \lambda_t$$

$$\lambda_t = \lambda_{t+1}$$

$$\lambda_T = \phi$$

$$\phi \geq 0 \text{ with } \phi W_{T+1} = 0 \Rightarrow \beta^T u_c(c_T) W_{T+1} = 0$$

$$u'(c_t) = \beta u'(c_{t+1}) \quad \forall t \in [0, T - 1]$$

$$W_{T+1} = 0$$

- With the set of T intertemporal equations (euler equations), an initial condition and a terminal condition

The Recursive Formulation

A Cake Eating Problem

- In order to solve finite-horizon dynamic programming problems, we are going to proceed by backwards induction.
- For $t = T$, given the properties of u and the constraint, the optimal solution is given by:

$$c_T = W_T$$
$$u(c_T) = u(W_T)$$

The Recursive Formulation

A Cake Eating Problem

- We define the **value function** at time T as:

$$V_T(W_T) = \max_{c_T} u(c_T)$$
$$c_T + W_{T+1} = W_T$$

- The optimal cake-saving decision is thus:

$$g_T(W_T) = 0$$

and the value function is given by:

$$V_T(W_T) = u(W_T)$$

The Recursive Formulation

A Cake Eating Problem

- Now let's go to $t = T - 1$ given that we have solved the problem for $t = T$ and define V_{T-1} .

$$\begin{aligned} V_{T-1}(W_{T-1}) &= \max_{c_{T-1}, c_T, W_T, W_{T+1}} u(c_{T-1}) + \beta u(c_T) \\ \text{s.t. } c_{T-1} + W_T &= W_{T-1} \\ c_T + W_{T+1} &= W_T \end{aligned}$$

- Given that we already we know what is optimal to do in the next period, we can simplify the problem at $T - 1$ as:

$$\begin{aligned} V_{T-1}(W_{T-1}) &= \max_{c_{T-1}, W_T} u(c_{T-1}) + \beta V_T(W_T) \\ \text{s.t. } c_{T-1} + W_T &= W_{T-1} \end{aligned}$$

The Recursive Formulation

A Cake Eating Problem

- Let's write the optimality conditions as:

$$u'(c_{T-1}) = \beta V'_T(W_T)$$

$$u'(c_{T-1}) = \beta u'_T(W_T)$$

- The solution coincides with the sequential formulation in the last period.
- We are in good track but what about previous periods?

The Recursive Formulation

A Cake Eating Problem

- Since it's going to be useful let's first derive the value of $V'_{T-1}(W_{T-1})$ given the optimal cake saving decision $g_{T-1}(W_{T-1})$ obtained from the previous FOC.

$$\begin{aligned}
 V_{T-1}(W_{T-1}) &= u(W_{T-1} - g_{T-1}(W_{T-1})) + \beta V_T(g_{T-1}(W_{T-1})) \\
 \frac{\partial V_{T-1}(W_{T-1})}{\partial W_{T-1}} &= u_c(c_{T-1}) - u_c(c_{T-1}) \frac{\partial g_{T-1}(W_{T-1})}{\partial W_{T-1}} + \\
 &\quad \beta \frac{\partial g_{T-1}(W_{T-1})}{\partial W_{T-1}} \frac{V_T(W_T)}{\partial W_T} \\
 \frac{\partial V_{T-1}(W_{T-1})}{\partial W_{T-1}} &= u_c(c_{T-1}) + \frac{\partial g_{T-1}(W_{T-1})}{\partial W_{T-1}} \left(\beta \frac{V_T(W_T)}{\partial W_T} - u_c(c_{T-1}) \right) \\
 \frac{\partial V_{T-1}(W_{T-1})}{\partial W_{T-1}} &= u_c(c_{T-1})
 \end{aligned}$$

The Recursive Formulation

A Cake Eating Problem

- At $T - 2$ the problem can be written as:

$$V_{T-2}(W_{T-2}) = \max_{c_{T-2}, W_{T-1}} u(c_{T-2}) + \beta V_{T-1}(W_{T-1})$$
$$\text{s.t. } c_{T-2} + W_{T-1} = W_{T-2}$$

- With FOCs:

$$u_c(c_{T-2}) = \beta \frac{\partial V_{T-1}(W_{T-1})}{\partial W_{T-1}} = \beta u_c(c_{T-1})$$

Infinite Horizon

A Cake Eating Problem

- Suppose for the cake-eating problem, we allow the horizon to go to infinity.
- The main advantage of an infinite horizon is that the agent problem becomes stationary: the maximization problem at date t is exactly the same as in period $t + 1$
- Unlike in finite horizon case, we don't have a terminal condition in the cake eating problem we will thus need to impose a transversality condition:

$$\lim_{t \rightarrow \infty} \beta^t u_c(c_t) W_{t+1} = 0$$

if discounted marginal utility is positive, the amount of cake needs to go to zero to rule out over-accumulation

Infinite Horizon

A Cake Eating Problem

- One can consider solving the infinite horizon sequence given by:

$$\begin{aligned} \max_{\{c_t, W_{t+1}\}_{t=0}^{\infty}} & \sum_{t=0}^{\infty} \beta^t u(c_t) \\ \text{s.t.} & c_t + W_{t+1} = W_t + y \quad \forall t \\ & \lim_{t \rightarrow \infty} \beta^t u_c(c_t) W_{t+1} = 0 \end{aligned}$$

- Written in recursive form:

$$V(W_t) = \max_{\{c_t, W_{t+1}\}} u(c_t) + \beta V(W_{t+1}) \quad (1)$$

$$\begin{aligned} \text{s.t.} & c_t + W_{t+1} = W_t + y \\ & \lim_{t \rightarrow \infty} \beta^t V(W_t) = 0 \end{aligned} \quad (2)$$

The transversality condition (2) is frequently avoided because assuming V being bounded, it is satisfied for $\beta < 1$.

Infinite Horizon

A Cake Eating Problem

- Equation (1) is referred as the Bellman equation.
- It is a functional equation: the unknown represents as function.
- By FOCs:

$$u_c(c_t) = \beta \frac{\partial V(W_{t+1})}{\partial W_{t+1}} \quad (3)$$

- Let's define $g(W)$ the optimal savings function associated with equation (1):

$$g(W_t) = \arg \max_{W_{t+1}} u(W_t + y - W_{t+1}) + \beta V(W_{t+1})$$

$$V(W_t) = u(W_t + y - g(W_t)) + \beta V(g(W_t))$$

Infinite Horizon

A Cake Eating Problem

- Provided that g is differentiable we can now compute:

$$\frac{\partial V(W_t)}{\partial W_t} = u_c(c_t) + \frac{\partial g(W_t)}{\partial W_t} \left(\beta \frac{\partial V(W_{t+1})}{\partial W_{t+1}} - u_c(c_t) \right)$$

$$\frac{\partial V(W_t)}{\partial W_t} = u_c(c_t) \Rightarrow \frac{\partial V(W_{t+1})}{\partial W_{t+1}} = u_c(c_{t+1})$$

- Then we can write equation (3) as:

$$u_c(c_t) = \beta u_c(c_{t+1})$$

Infinite Horizon

A Cake Eating Problem

- Under what conditions V exists? Is it unique?
- How to find V in the infinite horizon case?
- Is g a function or a correspondence? Is it differentiable?

The Dynamic Programming Approach

- Building on the intuition gained from the cake eating problem, we now consider a more formal treatment of the dynamic programming approach to answer the previous questions.
- We begin with the nonstochastic case and then add uncertainty to the formulation.

The Dynamic Programming Approach

- Consider the infinite horizon optimization problem of an agent with payoff function $\tilde{\sigma}(s_t, c_t)$.
- state vector: s_t ; control vector: c_t .
- Transition equation: $s_{t+1} = \tilde{\tau}(s_t, c_t)$.
- The state summarizes all the information from the past that is needed to make a forward-looking decision.
- $s \in \mathcal{S}$ and $c \in \mathcal{C}(s)$.
- Let β be the discount factor and assume $0 < \beta < 1$.

The Dynamic Programming Approach

- The sequential problem can be written as:

$$\begin{aligned} \max_{\{c_t\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} \beta^t \tilde{\sigma}(s_t, c_t) \\ \text{s.t. } s_{t+1} = \tilde{\tau}(s_t, c_t) \\ c_t \in \tilde{\mathcal{C}}(s_t) \end{aligned}$$

- We can rewrite the problem by imposing the law of motion of the state:

$$\begin{aligned} V^*(s_0) = \max_{\{s_{t+1}\}_{t=0}^{\infty}} \sum_{t=0}^{\infty} \beta^t \sigma(s_t, s_{t+1}) \\ \text{s.t. } s_{t+1} \in \mathcal{C}(s_t), \end{aligned}$$

Where V^* denotes the highest possible value the the objective function can reach

- The basic idea of dynamic programming is to turn the sequential problem into a functional equation:

$$V(s) = \max_{s' \in \mathcal{C}(s)} \sigma(s, s') + \beta V(s') \quad (4)$$

- Instead of choosing a sequence $\{s_t\}_{t=0}^{\infty}$, we choose a policy, which determines the control s' as a function of the state s .
- Given that V appears both in both sides of the equation 4 and thus it is defined recursively.
- Equation 4 is also referred as the Bellman equation after Richard Bellman, who was the first to introduce the dynamic programming formulation.
- A solution to the functional equation is thus a fixed point.

Math Review

Brouwer's Fixed Point Theorem

- Let \mathcal{M} be a nonempty compact (closed and bounded) convex set.
- Let T be a continuous function that maps each point $x \in \mathcal{M}$ to itself.
- Then T has a fixed point $x^* \in \mathcal{F}$ such that $T(x^*) = x^*$
- More questions:
 - To which set does V belong to?
 - Does the operator defined in the functional equation map each element of that set to itself?
 - Is the fixed point unique?

Math Review

What is a Contraction Mapping?

- Let (\mathcal{M}, d) be a metric space where \mathcal{M} is a set and d is a metric.

A metric space is a set and a function such that for all $x, y, z \in S$:

- $d(x, y) \geq 0$, with equality iff $x = y$
- $d(x, y) = d(y, x)$
- $d(x, y) \leq d(x, z) + d(z, y)$

- Let $T : \mathcal{M} \rightarrow \mathcal{M}$ be an function mapping \mathcal{M} into itself.
- If there exists a $\beta \in (0, 1)$ such that,

$$d(Tz_1, Tz_2) \leq \beta d(z_1, z_2) \quad \forall z_1, z_2 \in S$$

then T is a **contraction mapping** with modulus β .

- In other words, a contraction mapping brings elements of the space \mathcal{M} uniformly closer to one another.

Math Review

Contraction Mapping Theorem

Let (\mathcal{M}, d) be a complete metric space and suppose $T : \mathcal{M} \rightarrow \mathcal{M}$ is a contraction mapping.

A metric space is complete if every Cauchy sequence is a convergent sequence.

- A sequence $\{x_n\}_{n=0}^{\infty}$ is a Cauchy sequence if for all $\epsilon > 0$ there exists an $N \in \mathbb{N}$ such that for all $l, n > N$, $d(x_l, x_n) < \epsilon$
 - A sequence $\{x_n\}_{n=0}^{\infty}$ is a convergent sequence to $x_0 \in \mathcal{M}$ if for all $\epsilon > 0$, there exist here exists an $N \in \mathbb{N}$ such that for any $n > N$, $d(x_n, x_0) < \epsilon$
- Then, T has a **unique** fixed point \hat{z} and for any $z_0 \in \mathcal{M}$, and any $n \in \mathbb{N}$ we have $d(T^n z_0, \hat{z}) \leq \beta^n d(z_0, \hat{z})$.
 - That is there exists a unique $\hat{z} \in \mathcal{M}$ such that

$$T\hat{z} = \hat{z}$$

and regardless of the starting guess z_0 , the sequence $\{T^n z_0\}_{n=0}^{\infty}$ converges to \hat{z} .

Match Review

Blackwell's Sufficient Conditions for a Contraction

- Let $s \in \mathcal{S}$ and (\mathcal{M}, d) be the metric space where \mathcal{M} is the set of bounded function equipped with the sup norm.
- Let $T : \mathcal{M}(s) \rightarrow \mathcal{M}(s)$ satisfying:
 1. Monotonicity: If $W(s) \geq Q(s)$, for all $s \in \mathcal{S}$, then $TW(s) \geq TQ(s)$.
 2. Discounting: for any constant k there exists $\tilde{\beta} \in [0, 1)$ such that $T(W + k)(s) \leq T(W)(s) + \tilde{\beta}k$.
- Then T is a contraction.

Recursive Formulation

- In order to apply the Blackwell sufficient conditions, we need V to belong to the set of bounded functions.
- For this to be true, we need some assumption on the primitive objects.

Recursive Formulation

- $\sigma(s_t, s_{t+1})$ needs to be bounded so that it does not yield infinite returns: we cannot compare two choices of s_{t+1} that deliver infinite value.
- With $\beta \in (0, 1)$ and bounded σ , the V will be bounded for the problems that we will see in this course.
- Problems might arise in models of growth: you would need growth in the return function to be “smaller” than the rate of discounting such that discounted returns are bounded.
- This assumption will allow us to define the set of V : the set of continuous bounded functions.
- Equipped with the supremum norm forms a complete metric space.

Recursive Formulation

- If σ is continuous and \mathcal{C} is nonempty and compact (closed and bounded).

⇒ Unique value function satisfying the functional equation and therefore it is possible to find $V(x)$ by an iterative process

1. Select any initial value $V_0(s) \forall s \in \mathcal{S}$.
2. Define a sequence of functions:

$$V_n(x) = \max_{s' \in \mathcal{C}(s)} \sigma(s, s') + \beta V_{n-1}(s)$$

3. The sequence $\{V_0, V_1, \dots, V_n\}_{n=0}^{\infty}$ converges to V

Recursive Formulation

- Even if V is unique it could be that the policy associated could be a correspondence unless we put further restrictions of σ and \mathcal{C} :
 1. $\sigma(s, s')$: strictly concave, continuous, and differentiable.
 2. $\mathcal{C}(s)$: convex

⇒ We have a continuous and differentiable policy function

- The Enveloppe theorem holds:

$$\frac{\partial v(s)}{\partial s} = \frac{\partial \sigma(s, s')}{\partial s}$$

Is T a Contraction?

Blackwell's Sufficient Conditions: Monotonicity

- Let $Q(s) \leq W(s) \forall s \in \mathcal{S}$.
- Let $\phi_Q(s)$ be the policy function obtained from:

$$\phi_Q(s) = \arg \max_{s' \in \mathcal{C}(s)} \sigma(s, s') + \beta Q(s')$$

- Then,

$$\begin{aligned} TQ(s) &= \sigma(s, \phi_Q(s)) + \beta Q(\phi_Q(s)) \leq \sigma(s, \phi_Q(s)) + \beta W(\phi_Q(s)) \\ &= \max_{s' \in \mathcal{C}(s)} \sigma(s, s') + \beta W(s') = TW(s) \end{aligned}$$

Is T a Contraction?

Blackwell's Sufficient Conditions: Discounting

- This property is easy to verify in the dynamic programming problem:

$$\begin{aligned} T(W + k)(s) &= \max_{s' \in \mathcal{C}(s)} \sigma(s, s') + \beta(W(s') + k) \\ &= TW(s) + \beta k \end{aligned}$$

The Neoclassical Growth Model

- In 1928 Frank Ramsey, a young mathematician, posed the problem:
“How much of its income should a nation save?”
and developed a dynamic model to answer this question.
- Economic agent (a social planner) producing output from labor and capital who must decide how to split production between consumption and capital accumulation.

The Neoclassical Growth Model

The Planner's Problem

- Time is discrete.
- Production is given by $y_t = f(k_t)$ where k_t is capital. f satisfies inada conditions.
- The planner's problem is given by:

$$\begin{aligned} \max_{\{c_t\}_{t=0}^{\infty} \{k_{t+1}\}_{t=0}^{\infty}} & \sum_{t=0}^{\infty} \beta^t u(c_t) \\ \text{s.t.} & \quad c_t + k_{t+1} \leq f(k_t) + (1 - \delta)k_t \quad \forall t \end{aligned}$$

The Neoclassical Growth Model

The Planner's Problem

- Now let's write the planner's problem in recursive form:

$$V(k) = \max_{k' \in [0, f(k) + (1-\delta)k]} u\left(f(k) + (1-\delta)k - k'\right) + \beta V(k')$$

- The solution is characterized by:

$$u_c(c) = \beta \frac{\partial V(k')}{\partial k'} = \beta \left(\frac{\partial f(k')}{\partial k'} + 1 - \delta \right) u_c(c')$$

The Neoclassical Growth Model

The Planner's Problem

- In the one sector growth model we define the operator T to be:

$$TV(k) = \max_{k' \in [0, f(k) + (1-\delta)k]} \{u(f(k) + (1-\delta)k - k') + \beta V(k')\}$$

- We want to argue that this operator has as unique fixed point using the contraction mapping theorem.
- Thus we are going to do it using Blackwell's sufficient conditions.

The Neoclassical Growth Model

The Planner's Problem

- Monotonicity:

Let $\phi_Q(k) = \arg \max_{k' \in \Gamma(k)} u(f(k) + (1 - \delta)k - k' + \beta Q(k'))$

if $Q(k) \leq W(k)$, for all k

then $TQ(k) = u(f(k) + (1 - \delta)k - \phi_Q(k)) + \beta V(\phi_Q(k))$
 $\leq u(f(k) + (1 - \delta)k - \phi_Q(k)) + \beta W(\phi_Q(k)) \leq TW(k)$

- Discounting:

$$\begin{aligned} T(V + a)(k) &= \max_{k' \in \Gamma(k)} \{u(f(k) + (1 - \delta)k - k') + \beta(V(k') + a)\} \\ &= TV(k) + \beta a \end{aligned}$$

Solving the problem numerically

Discrete State Methods

- There exists a variety of numerical methods to solve dynamic programming problems like the Ramsey problem (projection, perturbation, parameterized expectation).
- The need of numerical methods arises from the fact that dynamic programming problems generally do not have tractable closed form solutions.
- Because of their simplicity, we are going to focus on discrete-state space methods.

Solving the problem numerically

Discrete State Methods

- In this case, the value function is a finite dimensional object.
- For instance, if the state space is one dimensional and has elements $\mathcal{S} = s_1, s_1, \dots, s_n$, the value function is just a vector of n elements where each element gives the value attained by the optimal policy if the initial state of the system is $s_n \in \mathcal{S}$.
- Drawback: curse of dimensionality.
 - ▶ If the the value function of an m -dimensional problem with n different points in each dimension is an array of n^m different elements and the computation time needed to search this array may be prohibitively high.

Solving the problem numerically

Value Function Iteration

- Given that Blackwell sufficient conditions hold, we can use the following pseudo-code for finding the value function:
 1. Make a guess for V_0 for all values of capital.
 2. Apply the operator T and recover $V_1 = TV_0$
 3. Compute distance between V_0 and V_1 .
 - 3.1 If V_1 and V_0 are close enough, stop.
 - 3.2 Otherwise set $V_0 = V_1$ and go back to 2.
- Once the algorithm has converged, you can simulate the path for capital of an economy with an initial capital endowment.

Solving the problem numerically

Value Function Iteration


- Define a grid with N points of capital between $[\underline{k}, \bar{k}]$ around the steady state level of capital.
- Define a value of V_0 for all the points in this grid. Let's say $V_0 = 0$ for all k .
- Given this V_0 , we can generate a vector for each level of capital k_i which elements are:

$$\begin{bmatrix} u(f(k_i) + (1 - \delta)k_i - k_1) + \beta V_0(k_1) \\ u(f(k_i) + (1 - \delta)k_i - k_2) + \beta V_0(k_2) \\ \vdots \\ u(f(k_i) + (1 - \delta)k_i - k_N) + \beta V_0(k_N) \end{bmatrix}$$

Solving the problem numerically

Value Function Iteration


- $TV_0(k)$ can be approximated by the maximum value of the elements of this vector.
- Looping through all values of $i \in [0, N]$ we will recover V_1 .
- Given V_1 we can recover V_2 .



Solving the problem numerically

Value Function Iteration

- We iterate until V_g and V_{g+1} are sufficiently close



Solving the problem numerically

Value Function Iteration



- Now that we have V , we need to recover $\pi(k)$ which is given by:

$$\pi(k) = \arg \max_{k'} \{u(k, k') + \beta V(k')\}$$

Solving the problem numerically

What do we aim for?


- A policy function:



Solving the problem numerically

Evolution of capital

- Given $\pi(k)$ we can simulate the transition towards the steady state for any $k_0 \in [\underline{k}, \bar{k}]$.



The NGM with Uncertainty

- Now, we focus on another application: the stochastic version of neoclassical growth model where shocks are going to affect firm's productivity.

$$y(z, k) = e^z f(k)$$

- We are going to assume that stochastic variables can take finitely many values.
- This restriction allows us to use Markov chains to represent uncertainty.

The Social Planner Problem

- The recursive formulation of this problem can be written as:

$$\begin{aligned} V(k, z) &= \max_{k'} \left\{ u(k, k', z) + \beta E_{z'|z} [V(k', z')] \right\} \\ \text{s.t. } k' &\in \Gamma(k, z) \equiv [0, e^z f(k) + (1 - \delta)k] \\ z' &= \rho z + \epsilon, \quad \epsilon \sim N(0, \sigma_\epsilon^2) \end{aligned}$$

Discretize an AR(1) using the Tauchen Method

- Method for discretizing an AR(1) process in N points.

$$z_t = \rho z_{t-1} + \epsilon, \epsilon \sim N(0, \sigma_\epsilon^2)$$

- Unconditional variance: $\sigma_z^2 = \frac{\sigma_\epsilon^2}{1 - \rho^2}$
- Create a (equally spaced) grid with first and last point in the grid q standard deviations away from the mean: $z_1 = -q\sigma_z, z_N = q\sigma_z$
 - ▶ Space between points: $dz = \frac{z_N - z_1}{N - 1}$
 - Fill the transition matrix Π :

$$\begin{aligned} \Pi(z_j|z_i) &= Pr \left[z_j - dz/2 \leq \rho z_i + \epsilon \leq z_j + dz/2 \right] \\ &= \Phi \left[\frac{z_j + dz/2 - \rho z_i}{\sigma_\epsilon} \right] - \Phi \left[\frac{z_j - dz/2 - \rho z_i}{\sigma_\epsilon} \right] \end{aligned}$$

Blackwell's sufficient conditions

- Let's define the operator T over function V^g as:

$$TV^g(k_t, z_t) = \max_{k_{t+1}} \left\{ u(k_t, k_{t+1}, z_t) + \beta \sum_{z_{t+1}} \Pi(z_{t+1}|z_t) V^g(k_{t+1}, z_{t+1}) \right\}$$

- Is operator T a contraction mapping? Are Blackwell's sufficient conditions satisfied?

Blackwell's Sufficient Conditions

- Monotonicity: If $V(k_t, z_t) \leq W(k_t, z_t)$ for all k_t and z_t :

$$\begin{aligned}
 TV(k_t, z_t) &= \max_{k_{t+1}} \{ u(k_t, k_{t+1}, z_t) + \beta \sum_{z_{t+1}} \Pi(z_{t+1}|z_t) V(k_{t+1}, z_{t+1}) \} \\
 &= u(k_t, g_v(k_t, z_t), z_t) + \beta \sum_{z_{t+1}} \Pi(z_{t+1}|z_t) V(g_v(k_t, z_t), z_{t+1}) \\
 &\leq u(k_t, g_v(k_t, z_t), z_t) + \beta \sum_{z_{t+1}} \Pi(z_{t+1}|z_t) W(g_v(k_t), z_{t+1}) \leq TW(k_t, z_t)
 \end{aligned}$$

- Discounting:

$$\begin{aligned}
 T[V(k_t, z_t) + a] &= \max_{k_{t+1}} \{ u(k_t, k_{t+1}, z_t) + \beta \sum_{z_{t+1}} \Pi(z_{t+1}|z_t) (V(k_{t+1}, z_{t+1}) + a) \} \\
 &= T[V(k_t, z_t)] + \beta a
 \end{aligned}$$



Discrete State-Space Methods: Value Function Iteration

- Make a guess of V^0 and loop over all combinations of capital and shocks and solve for:

$$V^1(k_i, z_j) = \max_{k' \in K} U(e^{z_j} f(k_i) + (1 - \delta)k_i - k') + \beta \sum_{z'} \Pi(z'|z_j) V^0(k', z')$$



- If V^1 and V^0 are different, then set $V^0 = V^1$ and iterate until convergence.

Policy function



Simulation

- Now we don't reach an steady state level of capital but a stationary distribution for capital.



The Social Planner

Characterizing the Solution

- Taking first-order conditions from recursive formulation we get:

$$\frac{\partial u(k, k', z)}{\partial k'} + \beta E_{z'|z} \frac{\partial V(k', z')}{\partial k'} = 0$$

$$u_c(k, k', z) = \beta E_{z'|z} \left[(e^{z'} f'(k') + 1 - \delta) u_c(k', k'', z') \right]$$